

BMR Grazing Blend



SORGO SORGHUM x
SUDANGRASS



- ✂ Improved digestibility and palatability, high stem sugar
- ✂ Small stem type; best for grazing
- ✂ Excellent recovery after grazing or cutting
- ✂ Prolonged grazing

The BMR grazing blend produces excellent grazing. With the different growth pattern of this blend you can count on grazing more days than a typical hybrid. The BMR trait in the hybrids ensures excellent palatability and digestibility. This blend produces a high level of sugar contributing to increase intake and energy. Begin grazing at 18 -24". Use rotational grazing and stocking rates of 2-3 head per acre under dryland or rainfed conditions.

AGRONOMIC TRAITS

Height:	60-80"
Maturity:	Varies
Regrowth:	Excellent
Midrib Type:	BMR
Plant Type:	Juicy Sweet
Photoperiod Sensitive:	No
Min./Max. pH:	6.0-7.5
Downy Mildew:	Resistant
Anthracoese:	Resistant

CROP USE INFORMATION

Ease of Establishment:	Good
Double Cropping:	Excellent
Dryland/Irrigated:	Excellent on Both
Hay/Baleage:	Excellent
Silage/Greenchop:	Will need to cure for silage moisture
Rotational Grazing:	Good
Continuous Grazing:	Excellent
Cover Crop:	Excellent
Digestibility:	Good IVTD, NDFD, TDN %
Palatability:	Sweet & Leafy
Fertilizer:	1-1¼ Lbs N per growing day/acre

SEEDING RATES

Seeds Per Pound:	16,000
Soil Temperature:	62°F
Seeding Depth:	1"-1.5"

Seeding Method	Harvest Stage	Dryland Lbs./Acre	Irrigated Lbs./Acre	Dryland Seed/Acre	Irrigated Seed/Acre
Drilled	Boot	25-40	40-55	400,000-640,000	640,000-880,000*
Broadcast	Boot	30-45	45-60	480,000-720,000	720,000-960,000

HARVEST

- BMR Grazing Blend can be harvested but should not be allowed to grow tall. Cut between 50-55 days or 5-6 feet
- When harvested, cut it in the boot stage and 6-8 inches above ground level for best regrowth
- Can be windrowed for grazing, hay or silage.
- Following a freeze, extreme drought, or fertilizer application followed by stress; remove the cattle from the field. See our guide for how to manage prussic acid and nitrates.